



MUHAMMAD AL-XORAZMIY NOMIDAGI
TOSHKENT AXBOROT TEXNOLOGIYALARI UNIVERSITETI
NUKUS FILIALI



«XALQ XO'JALIGI SOHASIDA ILG'OR TEXNOLOGIYALAR TADBIQI MUAMMOLARI»

MAVZUSIDAGI HUDUDIIY ILMIIY-TEXNIK KONFERENSIYASI

MA'RUZALAR TO'PLAMI



Chorvachilikda ilg'or texnologiyalar
va innovatsion yechimlar



Dasturlash, kiber xavfsizlik va qishloq
xo'jaligi fan sohalari integratsiyasi



Ta'lim va ishlab chiqarishda innovatsiyalar,
tahlil va prognozlash vositalari



27-dekabr 2023 yil

Konferensiya IL-392103072-
"Chorvachilik komplekslarini
elektron boshqarishning mobil
ilovasini yaratish" innovatsion
loyiha doirasida olib borilgan
ilmiy-amaliy tadqiqotlar
natijalariga bagishlangan



Nukus sh. A.Dosnazarov k. 74 uy



(61) 222-49-10



www.uzplf.uz



www.tatunf.uz

<i>B.Y.Geldibayev</i> Chorva komplekslarida sut mahsuldorligi haqidagi tahliliy hisobatlarni shakllantirishda kdd tahlil jarayonidan foydalanish	87
<i>G.G'.Artikova, M.Sh.Qazaqov</i> Xorazm viloyatida online chorva bozori qurish uchun mo'ljallangan mobil ilova tahlili.	91
<i>J.I.Dauletnazarov</i> Aqli dehqonchilikda foydalaniladigan texnologiyalar	94
<i>B.Y.Geldibayev</i> IoT qurilmalaridan ma'lumotlarni olish jarayoni tashkil etishda «Edge Computing»dan foydalanishning afzalliklari	98
<i>J.I.Dauletnazarov</i> IoTning qishloq xo'jaligida qo'llanilishi	100
<i>A.A.Temirov</i> IoT asosidagi aqli qishloq xo'jaligi uchun energiya tejamkor Edge-Fog-Cloud arxitekturasi	105
<i>D.A.Ernazarov</i> Qoramollarda oqsoqliklarni va tuyoq kasalliklarini erta aniqlash	109
<i>Э.С.Бабаджанов, Н.И.Калимбетов</i> Қорамол касалликларини С4.5 алгоритми орқали таснифлаш	113
II SHO'BA. DASTURLASH, KIBER XAVFSIZLIK VA QISHLOQ XO'JALIGI FAN SOHALAR INTEGRATSIYASI	117
<i>A.X.Nishanov, B.C.Samandarov</i> Real vaqt rejimida dinamik ma'lumotlar oqimini samarali boshqarish masalasi	117
<i>A.X.Nishanov, X.B.Kenjaev</i> Matnlarni kalit so'zlar asosida umumlashtiruvchi tizimni yaratish vazifalari	121
<i>N.U.Uteuliev, G.M.Djaykov, D.Sh.Yuldoshev</i> Numerical method for solving the problem of integral geometry on a family of semicircles	123
<i>X.N.Zaynidinov, X.Sh.Quzibayev</i> Sun'iy neyron tarmoq yordamida quyi amudaryo hududidagi suv sifatini bashoratlash	127
<i>B.B.Akbaraliyev, R.X.Xoliqnazarov</i> Tashkilotlarga ichki elektron hujjat aylanuv tizimini joriy etish	131
<i>Sh.R.G'ulomov</i> Uzfirwall-Next Generation Firewall apparat-dasturiy vositasining funksional strukturasi	136
<i>T.T.Berdimbetov, S.K.Nietullayeva, G.Q.Baytileuova, D.O.Madetov, M.J.Eshbayev</i> GIS ilovalarining rivojlanish tendensiyalari	140
<i>T.T.Berdimbetov, S.K.Nietullayeva, G.Q.Baytileuova, D.O.Madetov, M.J.Eshbayev</i> GISta fazoviy mal'umotlar tahlili	143
<i>F.K.Achilova</i> "Hand Tools" mobil ilovasini ishlab chiqish va tadbiq etishning afzalliklari	146
<i>M.E.Shukurova</i> Neft qatlamlari g'ovak muhitida filtratsiya jarayoni chegaraviy masalalarini yechishni avtomatlashtirish	150
<i>D.Kenjaboeva</i> Ta'lim berishda o'qituvchi deontologisi va kompetentligi	154
<i>A.M.Risnazarov</i> Kishi resursli kriptografiya	157
<i>S.X.Saparov, U.B.Allayarov, H.B.Qudratov</i> Bosh miya saratoni kasalligini erta tasniflashda informativ belgilar majmuasini tanlash algoritmi	159
<i>S.X.Saparov, U.B.Allayarov, H.B.Qudratov</i> Bosh miya saratonini erta tasniflashda obyektlar muhimligini aniqlash algoritmi	164

- To‘plam hujjatlarini avtomatik umumlashtirish dasturiy vositasining infratuzilmasi va unda axborot oqimimini boshqarishning IDEF modellari;
- Dasturiy vosita arxitekturasi, ishlash mexanizimi va uni ishlab chiqish.

Bu vazifalar bajarilishi bo‘yicha juda uzviylikni tashkil etishi bilan birga ayniqsa 1-4 bo‘g‘inlar o‘zaro ajralmas va bir-birini to‘ldiruvchi asosi hisoblanadi. Endi mazkur vazifalarning mohiyatiga va ularni yechish usullari yoki mexanizmlari batafsilroq to‘xtalinadi.

ARS-Uz tizimi uchun qo‘yolgan vazifalar, keltirilgan ob‘ektlari va ulardagi munosabatlari bo‘yicha uning infratuzilmasini 1-rasmdagi kabi taklif etiladi.

ARS-Uz tizimi infratuzilmasini tashkil etuvchi har bir elementi alohida vazifa hisoblanib, ularni 4 qismga ajratish mumkin: maxsus ma’lumotlar va bilimlar bazasi; matematik-algoritmik ta’minot; kiruvchi-chiquvchi hujjat va foydalanuvchi interfeyslari. Endi tizim asosi bo‘lgan MB loyihalash masalasi qaraladi.

Maqolada doimiy rivojlanishda bo‘lgan matnli hujjatlarni avtomatik qayta ishlash yo‘nalishlari hisoblangan kalit so‘zlarni ajratib olish, matnni umumlashtirish va nomlangan ob'ektni tanish uslubiyatlari tadqiqatlari natijasida o‘zbek tilida berilgan bir jinsli elektronn matnli hujjatlarni predmet sohalari asosida umumlashtirish masalasi va uni yechish vazifalari keltirildi.

Foydalanilgan adabiyotlar

1. Nishnov A.X., Babadjanov E.S., Kenjaev X.B. Avtomatik matnlarni umumlashtirish usullari tahlili //Journal of Advances in Engineering Technology. – 2023. – №. 2. – C. 37-46.
2. Kenjaev X.B. Hujjatlardan kalit so‘zlarni chiqarish algoritmlari va matn o‘xshashligini baholash usullari tahlili // Muhammad al-Xorazmiy avlodlari 3(25)/2023. b. 39-44
3. A.X.Nishanov, X.B.Kenjayeov, Hujjatlardan jadvallarni chiqarib olish masalasi, usullari va dasturiy ta’minotlar tahlili // Digital Transformation and Artificial Intelligence, ISSN: 3128-8121. Vol 1, No.2. 2023

NUMERICAL METHOD FOR SOLVING THE PROBLEM OF INTEGRAL GEOMETRY ON A FAMILY OF SEMICIRCLES

f-m.f.d., prof. N.U.Uteuliev (Nukus branch of TUIT),

PhD G.M.Djaykov (Nukus branch of TUIT),

D.Sh.Yuldoshev (Nukus branch of TUIT)

Abstract. We study the problem of integral geometry on a family of semicircles. A numerical algorithm for finding an approximate solution based on Tikhonov regularization is constructed.

Keywords: Integral geometry problems, inversion formula, algorithm, regularization, Radon transform

Reconstructing a function based on its spherical Radon transform with limited integration centers forms the foundation of several modern visualization technologies. This approach is employed in various fields, ultrasound imaging, Synthetic-aperture radar, as well as photoacoustic and thermoacoustic tomography.

Motivated by applications in thermoacoustic tomography, the problem of recovering a function supported in a ball from its average values over spheres centered on the boundary of the ball has received considerable attention in the last decade, with results appearing in various papers during this decade [1, 2].

Let's denote $L_H = \{(x, y) : x \in R^1, 0 \leq y \leq H\}$. For everyone (x, y) lying in the lane L_H . Let us denote by $C_0^2(L_H)$ the class of functions $u(x, y)$ that have all continuous partial derivatives up to 2 – order inclusive and are compactly supported with support in the strip L_H .

Statement of the problem. In a strip L_H restore a function of two variables $u(x, y)$ if the integrals of it over the curves of the family are known $\{\Upsilon(x, y)\}$

$$\int_{\Upsilon(x,y)} g(x, \xi) u(\xi, \eta) d\xi = f(x, y) \quad (1)$$

where an arbitrary family curve is represented by the expression

$$\Upsilon(x, y) = \left\{ (\xi, \eta) : (x - \xi)^2 + \eta^2 = y^2, 0 \leq \eta \leq y \leq H, -\infty < x < \infty \right\}.$$

When restoring function, $u(x, y)$ the following questions arise? 1. Uniqueness recovery function $u(x, y)$; Inversion formulas and recovery algorithms, recovery stability.

All of these questions are essentially answered for the classical Radon transform occurring in X-ray CT, positron emission tomography, and magnetic resonance imaging. However, they are much more complex and not as well understood for the circular Radon transform occurring in the thermoacoustic tomography.

1. Inversion formula

Let us rewrite equation (1) for $g(x, \xi) = 1$ in the following form

$$\int_{\Upsilon(x,y)} u(\xi, \eta) d\xi = f(x, y) \quad (2)$$

The solution to equation (2) has the following form:

$$\hat{u}(\lambda, y) = \frac{1}{\pi y} \frac{\partial}{\partial y} \int_0^y \frac{\eta ch\left(\lambda \sqrt{y^2 - \eta^2}\right)}{\sqrt{y^2 - \eta^2}} \cdot \hat{f}(\lambda, \eta) d\eta.$$

2. Sequence of approximate solutions integral geometry problems based on the regularizing operator

Firstly, in the last formula you can notice that for a fixed $0 \leq \eta \leq y \leq H$ $ch(\lambda\sqrt{y^2 - \eta^2})$ tend to infinity at $|\lambda| \rightarrow \infty$.

Secondly, instead of $f(\lambda, \cdot)$ measured $f^\delta(\lambda, \cdot) = f(\lambda, \cdot) + \delta f(\lambda, \cdot)$, where is $\delta f(\lambda, \cdot)$ the measurement error. The Fourier transform can be written as

$$f^\delta(\lambda, \cdot) = \int_{-\infty}^{+\infty} f(x, \cdot) e^{i\lambda x} dx + \int_{-\infty}^{+\infty} \delta f(x, \cdot) e^{i\lambda x} dx$$

where $\int_{-\infty}^{+\infty} \delta f(x, \cdot) e^{i\lambda x} dx$ is the Fourier transform of the noise, which usually

contains a white noise component. Since when $\lambda \rightarrow \infty$ $\int_{-\infty}^{+\infty} \delta f(x, \cdot) e^{i\lambda x} dx \rightarrow const$ and the integral $f^\delta(\lambda, \cdot)$ diverges.

If we want to build approximate solutions for the last equation that are stable to small deviations of the right-hand side $f(x, y)$, using the inverse Fourier transform, then we need to suppress the influence of high frequencies λ .

Consider the operator

$$R(u, \alpha) = L^{-1} \left[\psi(\hat{u}(\lambda, y), \lambda) \cdot W(\lambda, \alpha) \right]$$

where L^{-1} is the inverse Fourier transform, and $W(\lambda, \alpha)$ – is a certain given function defined for all non-negative values of the parameter α and any λ for which the inverse Fourier transform is taken.

Let the function $W(\lambda, \alpha)$ satisfy the following conditions:

1. $W(\lambda, \alpha)$ defined in area $(\alpha \geq 0, -\infty < \lambda < +\infty)$;
2. $0 \leq W(\lambda, \alpha) \leq 1$ for all values $\alpha \geq 0$ and λ ;
3. $W(\lambda, 0) \equiv 1$;
4. for everyone $\alpha > 0$ $W(\lambda, \alpha)$ – even in λ and $W(\lambda, \alpha) \in L_2(-\infty, \infty)$;
5. for everyone $\alpha > 0$ $W(\lambda, \alpha) \rightarrow 0$ at $\lambda \rightarrow \pm\infty$;
6. for $\alpha \rightarrow 0$ $W(\lambda, \alpha) \rightarrow 1$, without decreasing, and on any segment $|\lambda| \leq \lambda_1$ this convergence is uniform;
7. for everyone $\alpha > 0$ $ch(\lambda \cdot) W(\lambda, \alpha) \in L_2(-\infty, \infty)$;

8. for any $\lambda \neq 0$ $W(\lambda, \alpha) \rightarrow 0$ and $\alpha \rightarrow 0$ this convergence is uniform on any segment $[\lambda_1, \lambda_2]$, $0 < \lambda_1 < \lambda_2$.

If the function $W(\lambda, \alpha)$ satisfies conditions 1-8, then the operator defined with its help $R(u, \alpha)$

$$R(u, \alpha) = \frac{1}{\pi y} \frac{\partial}{\partial y} \int_0^{y+\infty} \int_{-\infty}^y \frac{\eta ch(\lambda \sqrt{y^2 - \eta^2}) W(\lambda, \alpha)}{\sqrt{y^2 - \eta^2}} \cdot \hat{f}(\lambda, \eta) d\eta$$

is the regularizing operator for equation (2) [3].

Let's consider a method for constructing a regularizing operator in the following form

$$u_{\alpha_1}(x, y) = \frac{1}{\pi y} \frac{\partial}{\partial y} \int_0^{y+\infty} \int_{-\infty}^y \frac{ch^2(\lambda \sqrt{y^2 - \eta^2})}{ch^2(\lambda \sqrt{y^2 - \eta^2}) + \alpha_1(1 + \lambda^2)} \cdot \frac{\eta \hat{f}(\lambda, \eta) e^{-i\lambda x}}{\sqrt{y^2 - \eta^2}} d\lambda d\eta$$

3. Numerical experiment

For the numerical experiment, we introduce uniform grids along x, y , $x \in [-a, a], y \in [c, d]$. $x_k = -a + kh_x$, $y_j = c + jh_y$, Where $h_x = \frac{2a}{n_x}, h_y = \frac{d-c}{n_y}$.

The discrete Fourier transform for a function $f(x_i, \cdot)$ is determined by the formula

$$\hat{f}(\lambda_m, \cdot) = \sum_{k=0}^{n_x-1} f(x_k, \cdot) e^{i\lambda_m x} = \sum_{k=0}^{n_x-1} f(x_k, \cdot) e^{i2\pi km/n_x}, m = 0, 1, \dots, n_x - 1$$

Here $\lambda_m = m\Delta\lambda$, $\Delta\lambda = \frac{2\pi}{T}$.

Inverse discrete Fourier transform

$$f(x_k, \cdot) = \sum_{m=0}^{n_x-1} \hat{f}(\lambda_m, \cdot) e^{-i2\pi km/n_x}, k = 0, 1, \dots, n_x - 1$$

Taking into account the above, we rewrite the inversion formula in the following form:

$$u_{\alpha_2}(x_k, y_j) = \frac{1}{\pi y} \frac{\partial}{\partial y} \int_0^{y_j} \frac{T(x_k, y_j, \eta) d\eta}{\sqrt{y_j^2 - \eta^2}} \tag{3}$$

Where $T(x_k, y_j, \eta) = \int_{-a}^{+a} \frac{ch^2(\lambda \sqrt{y_j^2 - \eta^2}) \eta \hat{f}(\lambda, \eta) e^{-i\lambda x_k}}{ch^2(\lambda \sqrt{y_j^2 - \eta^2}) + \alpha_1(1 + \lambda^2)} d\lambda$

After calculating the values of the functions, $T(x_k, y_j, \eta)$ integral (3) is calculated using non-standard quadrature formulas, which explicitly take into account the nature of the singularity.

We use the following function as a test example

$$u(x, y) = (x^2 - 1) \left(y^2 - \frac{y}{2} \right)$$

We present the results for the exact right-hand side in Fig.1 and, for comparison, show the results using formula (3).




Fig.1. Results of restoring the test function. a) exact solution, b) restoration according to the formula

References

1. Uteuliev N.U., Djaykov G.M., Seidulleev A.K. Inversion formula for the problem of integral geometry on families of parabolas. AIP Conference Proceedings July 2021. 2365(1):070004. DOI: [10.1063/5.0056852](https://doi.org/10.1063/5.0056852)
2. Begmatov A.Kh., Djaykov G.M. On the restoration of a function from spherical means. // Reports of the Academy of Sciences of the Higher School of the Russian Federation. 2013. Vol.1. No. 20. P. 6-16.
3. Tikhonov A.N., Arsenin V. Solutions of ill-posed problems. – 1977.

SUN'IY NEYRON TARMOQ YORDAMIDA QUYI AMUDARYO HUDUDIDAGI SUV SIFATINI BASHORATLASH

t.f.d., prof. X.N.Zaynidinov (TATU)

X.Sh.Quzibayev (TATU tayanch doktoronti)

Annotatsiya. Suv havzalaridagi fizik, kimyoviy va biologik jarayonlarning murakkab muhitida suv sifati o'zgaruvchilarining chiziqli bo'lmagan xatti-harakatlarini hisobga olgan holda suv sifatinii modellashtirish qiyin. Quyii Amudaryo